


Body Features Used in Identification



Please refer to your State or Provincial fishing guidelines for creel and size regulations on each species.


For additional information, please contact:
 Stacy Furgal, Fisheries and Ecosystem Health Specialist
 315-312-3042
 slf85@cornell.edu

This brochure results from project M/PIU-12 funded under award NA07OAR4170010 from NOAA's National Sea Grant College Program to the Research Fdn of SUNY on behalf of New York Sea Grant. Funding for the printing of this document was provided by the New York State Environmental Protection Fund under the authority of the New York Ocean and Great Lakes Ecosystem Conservation Act.


All images in this publication are property of New York Sea Grant and are not to be duplicated or published without permission.

Atlantic Salmon


(*Salmo salar*)
 Lake Phase Adult



Stream Phase Adult




Large kypes (jaw "hooks") on spawning males




Spawning Atlantic Salmon and Brown Trout can be difficult to distinguish. Vomerine teeth are the best characteristic to distinguish between these species.

Juvenile (parr stage)




Brown Trout


(*Salmo trutta*)
 Lake Phase Adult



Stream Phase Adult




Large kypes (jaw "hooks") on spawning males



Spawning Atlantic Salmon and Brown Trout can be difficult to distinguish. Vomerine teeth are the best characteristic to distinguish between these species.

Juvenile (parr stage)



Salmon and Trout of the Great Lakes:

A Visual Identification Guide




Original By: Dave MacNeill, Mary Austerman (NY Sea Grant)
 Dan Bishop, Fran Verdoliva (NYSDEC)
 Jim Johnson (USGS)
 Revised By: Jesse Lepak (NY Sea Grant)
 Dan O'Keefe (MI Sea Grant)
 Mitchell Zischke (IL-IN Sea Grant)
 Titus Seilheimer (WI Sea Grant)
 Artwork By: Peter Thompson




Chinook Salmon

(*Oncorhynchus tshawytscha*)


Lake Phase Adult



Stream Phase Adult




Large kypes (jaw "hooks") on spawning males



Pink Salmon and Chinook Salmon can hybridize. These hybrids are called "Pinooks."

Pinooks have variable characteristics and their appearance can be similar to either parent species.


Juvenile (parr stage)




Coho Salmon

(*Oncorhynchus kisutch*)

Lake Phase Adult




Early Stream Phase Adult



Late Stream Phase Adult




Juvenile (parr stage)




Steelhead/Rainbow Trout

(*Oncorhynchus mykiss*)


Lake Phase Adult



Stream Phase Adult



Large kypes (jaw "hooks") on spawning males




Spawning fish reddish to pink stripe

Lake Trout and Brook Trout can hybridize. These hybrids are called "Splake."

Splake have variable characteristics and their appearance can be similar to either parent species.

Juvenile (parr stage)



Pink Salmon

Lake Superior and St. Mary's River
Rare in Western Great Lakes

(*Oncorhynchus gorbuscha*)

Lake Phase Adult




Stream Phase Adult



Lake Trout


(*Salvelinus namaycush*)



Brook Trout

Lake Superior and streams

(*Salvelinus fontinalis*)



Fish coloration varies greatly and live fish may not look exactly as shown in the brochure.