

Geology

The Great Lakes were formed by the advancing and retreating of glaciers that gouged and shaped the basin over thousands of years from the great ice Age to approximately 10,000 years ago.

When the lakes took on their present shape. Over several different glacial periods, ancient rivers provided the beginnings of some of the Great Lakes. The east-west fetch of lake Erie followed one

until it reached its present size and shape. Like giant bulldozers, the of these ancient rivers and became wider and deeper over time.

power for lumber and grainmills), further impacting the fishery as and Atlantic salmon. Early settlements built dams to harness water

resulted in overfishings of important species like whitefish

Commission was established by the Canadian-U.S. Convention on the Great Lakes Fishery Commission coordinates fisheries research, facilitates cooperative fishery management among the state, provincial, tribal, and federal management agencies, and is responsible for controlling the invasive sea lamprey.

Modern Great Lakes

Pollution soon became a problem around the great lakes, as more and more people crowded cities in search of jobs and a better life than they left in Europe. Human and industrial wastes found their way to the once pristine waters of the Great Lakes as contaminants, heavy metals and industrial by-products created water quality issues. Phosphorus level increased and brought about a growth of nuisance algae. Areas of low oxygen levels (anoxia) caused fish die offs and in the late 1960's Lake Erie was considered "dead". An outcry from concerned citizens soon brought about a ban on detergents that contained phosphorus and other efforts to improve the water quality of the Great Lakes.

These early environmental efforts helped to pass the **Great Lakes Water Quality Agreement** between Canada and the U.S. in 1972. The Agreement was amended in 1987, and most recently in 2012, when provisions were added to address aquatic invasive species, habitat degradation, and the effects of climate change.

INVASIVE SPECIES:

From the spread of the sea lamprey with the opening of the Erie and Welland Canals in the early 1980's, invasive species have impacted the environment of the Great Lakes. Today, more than 180 invasive species, from the alewife to zebra mussels, make their home in the Great Lakes since their arrival in ships' ballast water in the late 1980s. The original zebra mussel (*Dreissena polymorpha*) and the related quagga mussel (*Dreissena rostriformis bugensis*) that arrived a few years later, are responsible for changes in the food web. They filter out plankton and other particles from the water, reducing the available food for other species. Both species of mussels are prolific breeders and their numbers grew rapidly. Quagga mussels are found in colder, deeper areas of the Great lakes, often covering the substrate hundreds of feet below the surface.

Zeبرا and quagga mussels have had a profound impact on the Great Lakes and have caused the extinction of some native species.

Atmospheric depositions also pose a threat to the health of the Great Lakes Basin. These chemicals come from industrial smokestacks, coal-burning power plants, automobile emissions and the spraying of pesticides. Although the United States and Canada have enacted environmental protection laws and regulations, the airshed of the Great Lakes can be impacted by chemicals that originate from areas like Mexico and Central America, where laws are not as stringent. The chemicals in the air are washed back down to land through precipitation, in the form of rain and snow.

Today, it is the emerging chemicals of concern that are the focus of research and study. Fire retardants that are put on clothes and furniture, pharmaceutical byproducts that are flushed down toilets or poured down drains, microplastics that can be found in body wash and toothpaste all have the potential to harm the environment. Some of the chemicals are endocrine disruptors that have been reported as causing reproductive issues in animals and fish.

Although many aquatic invasive species (AIS) have spread through the Great Lakes by ballast water, bait bucket dumping and hitchhiking on recreational boats and trailers, these are not the only means of introductions. Recently, efforts have been made to prevent the introduction of AIS through the aquarium and nursery trades and classroom releases. These **organisms in trade (OIT)** pathways provide a significant threat to the health of

H	— Huron
O	— Ontario
M	— Michigan
E	— Erie
S	— Superior

A good way to remember the names of the Great Lakes is by using the word "HOMES" as an acronym:

New York Sea Grant
229 Jarvis Hall
SUNY at Buffalo
Buffalo, NY 14260-4400
Tel: 716.645.3610

www.nyseagrant.org

New York's Sea Grant Extension Program provides Equal Program and Equal Employment Opportunities in association with Cornell Cooperative Extension, U.S. Department of Agriculture and U.S. Department of Commerce and cooperating County Cooperative Extension Associations.
--

Although a water restriction crisis in the Toledo, Ohio, area during the summer of 2014 focused the national spotlight on this issue, harmful algal blooms are not a new phenomenon to Lake Erie. During the 1960s and 1970s, the lake had massive cyanobacteria (blue-green algae) blooms due to cultural eutrophication. As the lake was cleaned up, these blooms seemed to diminish until the mid to late 1990s. Since then, the lake has seen significant numbers of HABs and recent blooms have been much more widespread than those of the past.

Habitat Destruction

Habitat destruction is a major issue in the Great Lakes, especially the ecologically important wetland areas that serve as spawning and nesting habitat for fishes and animals. Wetlands are some of the most biologically diverse areas in the basin and play a major role in water quality and erosion control. Unfortunately, many of the wetlands of the Great Lakes have been filled in and destroyed. In some areas of the Great Lakes, up to 90% of the original wetland areas are gone, being replaced by agriculture, industry or shoreline communities.

Land use practices continue to be a concern throughout the Great Lakes Basin with development and urban sprawl changing the use of the land and water drainage from natural to altered states. The functioning of natural systems and the connectivity of habitats can be forever changed without proper planning. Communities need to focus on land use and sustainable planning to reduce the impact on the basin. A balance needs to be created between development and environmental protection, with the goal of conserving and enhancing natural areas as part of land use planning.

Pollution

Early on, the primary threat from pollution came from unregulated industrial dumping and discharges from antiquated water treatment facilities. This type of pollution is referred to as **point source pollution** since it comes from an identifiable source like the end of a discharge pipe. Contaminants like PCBs, mercury, mirex, and other pollutants impact the food web as they bioaccumulate in the tissues of birds, fish and mammals and move from one trophic level to another. In the past, chemical concentrations were so high that fish consumption advisories were put in place to protect residents around the basin. Even today, restrictions are still in place for some species of bottom-dwelling fish like bullhead and carp.

Fortunately, efforts have been made to strengthen or enact environmental laws and regulations to limit point source pollution in the Great Lakes, and millions of dollars have been spent to upgrade or replace outdated water treatment plants around the basin.

Nonpoint pollution comes from many diffuse sources and it is much harder to identify the point of origin. The primary source of nonpoint source pollution is runoff. Development and growth mean more roads, parking lots, roofs and other impervious surfaces that allow for the runoff of chemicals, fertilizers, oil and gasoline into streams, lakes and other waterways. Phosphorus and nitrogen run off residential lawns, parks, golf courses and farms, causing eutrophication problems in the Great Lakes.

Sea Grant
New York

New York's Sea Grant Extension Program provides Equal Program and Equal Employment Opportunities in association with Cornell Cooperative Extension, U.S. Department of Agriculture and U.S. Department of Commerce and cooperating County Cooperative Extension Associations.
--

in many winters. The coastal regions of the Great Lakes contain microclimate areas that are less prone to late spring and early fall frosts, which is ideal for the many fruit orchards and vineyards that add to the economy of the basin.

The Great Lakes are an incredible resource that influences the lives, economies and communities that ring their shoreline. From tourism that brings millions of visitors to parks and beaches, to shipping that utilizes this "Highway H₂O" to move cargo, the Great Lakes serve as an economic driver to North America. The importance of the Great Lakes is not only counted in dollars and cents, but also the intrinsic value of their beauty and majesty by those who take in their scenic views and natural lands.

young, fledgling America become a world leader.

and their boundless resources played an important role in helping resources like fish and animals from the region. The Great Lakes as people cleared the land for agriculture and relied on the natural available food for other species. Both species of mussels are prolific breeders and their numbers grew rapidly. Quagga mussels are

the Midwest. Soon, towns and cities spread out across the basin and goods to the expanding regions around the Great Lakes and shores of America. The Erie Canal also provided a way to move peo-

the white pine and other lumber of the Great Lakes to the eastern The opening of the **Erie Canal** in 1825, brought a new way to ship Great Lakes region and European settlement steadily increased.

modify. The British soon realized the resources available from the trade in furs, especially beaver pelts, became an important com- ing the harsh conditions. The French continued to settle the basin as the Native Americans, they established settlements and survived in Orient through the Northwest Passage. Instead, with the help of Lakes. These settlers came in search of a new passage to the Great Early European settlers are given credit for "discovering" the Great

(Chippewa), Fox, Iroquois, Ottawa, Potawatomi and other tribes.

Over time, early Native Americans in the region included Ojibwe early settlers who were able to use copper to form tools and weapons. The Paleo-Indians were followed by as the Great Lakes Basin. These hunters depended on mammals like The Paleo-Indians were the first people to occupy the area we know

Early History

corals and large trilobites can be unearthed around the basin.

today, fossils from prehistoric sea creatures like clams and crinoids, the Great Lakes as deposits of halite, gypsum, oil and gases. Even sea. The sands, salts and minerals of that sea can be found beneath area where the basin lies was once covered by a shallow, tropical Millions of years before the glaciers carved out the Great Lakes, the

as the "flowerpots" on Bruce Peninsula in Ontario, Canada.

huge boulders and unusual rock formations that can be seen today ara Falls. These glaciers were more than a mile thick and left behind York, Michigan, Illinois and Wisconsin), and formed the mighty Niagara Escarpment that covers parts of Ontario, New sandstone and shale. As the glaciers moved over the area, they glaciers moved the resistant bedrock and easily scoured the softer until it reached its present size and shape. Like giant bulldozers, the of these ancient rivers and became wider and deeper over time. power for lumber and grainmills), further impacting the fishery as and Atlantic salmon. Early settlements built dams to harness water resulted in overfishings of important species like whitefish

As cities grew, so did the inevitable toll on the natural resources of the Great Lakes. Commercial fishing played an important role in the

water transparency in the Great Lakes. The increased light filtering activity of zebra and quagga mussels has increased

consumed by bottom-dwelling fish like round gobies.

bacteria that can become part of the food web when mussels are of the lakes, they can actually take up pollutants, contaminants and the water clearer. In fact, as mussels filter water close to the bottom

from the black and Caspian Seas in Europe. These aggressive

Like zebra and quagga mussels, the **round goby** (*Neogobius melanostomus*) was introduced in ballast water from ships coming

numbered. Gobies are easily identified by their suction cup-shaped, fused pelvic fins. No other Great Lakes fish has this suction disk

on its belly. Round gobies have highly developed sensory systems that allow them to avoid predators and detect prey. They are also capable of multiple spawnings (up to 6 times per year), allowing

them to quickly increase and maintain the size of their populations.

The **sea lamprey** (*Petromyzon marinus*) is referred to as the "vampire" of the Great Lakes since it is a parasitic fish that feeds

with the opening of the Erie and Welland Canals. Sea lampreys have a sucker-like mouth lined with sharp teeth that they use to attach to fish. Once firmly anchored, sea lampreys use a rasping tongue to break through the slime and scales before feeding. Special enzymes are released by the lamprey to continue its meal. Sea lampreys take a toll on the fishes of the Great Lakes, often targeting fish with small scales like whitefish, lake trout and salmon. The Great Lakes Fishery Commission is charged with controlling sea lamprey populations and they use a variety of techniques including Lampricides (pesticides to target lampreys), traps and barriers. People are encouraged to help reduce the spread of invasive species by cleaning their boats and trailers before moving them to other areas, draining live-wells, properly disposing of bait and taking other steps. Visit <http://www.protectyourwaters.net/> to learn more about the Stop Aquatic Hitchhiker! campaign and what you can do to slow the spread of aquatic invasive species.

Although many aquatic invasive species (AIS) have spread through the Great Lakes by ballast water, bait bucket dumping and hitchhiking on recreational boats and trailers, these are not the only means of introductions. Recently, efforts have been made to prevent the introduction of AIS through the aquarium and nursery trades and classroom releases. These **organisms in trade (OIT)** pathways provide a significant threat to the health of

Habitat destruction is a major issue in the Great Lakes, especially the ecologically important wetland areas that serve as spawning and nesting habitat for fishes and animals. Wetlands are some of the most biologically diverse areas in the basin and play a major role in water quality and erosion control. Unfortunately, many of the wetlands of the Great Lakes have been filled in and destroyed. In some areas of the Great Lakes, up to 90% of the original wetland areas are gone, being replaced by agriculture, industry or shoreline communities.

Great Lakes Basin Map



www.nyseagrant.org

